CORRECTION Open Access

Correction: Measurement properties of PROMIS short forms for pain and function in patients receiving knee arthroplasty



Anika Stephan^{1*}, Vincent A. Stadelmann¹, Stefan Preiss² and Franco M. Impellizzeri^{1,3}

Correction: Journal of Patient-Reported Outcomes (2023) 7:18

https://doi.org/10.1186/s41687-023-00559-x

Following publication of the original article [1], the authors identified that the original colours did not display correctly in Table 2. The correct table is given below.


The original article [1] has been updated.

The original article can be found online at https://doi.org/10.1186/s41687-023-00559-x.

*Correspondence: Anika Stephan anika.stephan@kws.ch

¹ Department of Teaching, Research and Development – Lower Extremities, Schulthess Clinic, Lengghalde 2, 8008 Zurich, Switzerland

³ Faculty of Health, University of Technology Sydney, PO Box 123, Broadway, NSW 2007, Australia



² Knee Surgery, Schulthess Clinic, Lengghalde 2, 8008 Zurich, Switzerland

Table 2 Correlations between PROMIS scales and the OKS

		Correlation with OKS		Hypothesis testing group
PROMIS PAIN	Baseline	Total:	- 0.69 (- 0.77 to - 0.58) ^a	Construct validity
		Men:	- 0.65 (- 0.78 to - 0.47) ^b	Construct validity
		Women:	- 0.72 (- 0.82 to - 0.60) ^a	Construct validity
	12 months	Total:	- 0.69 (- 0.78 to - 0.57) ^a	Construct validity
		Men:	- 0.64 (- 0.77 to - 0.45) ^a	Construct validity
		Women:	- 0.71 (- 0.84 to - 0.53) ^a	Construct validity
	Change	Total:	- 0.66 (- 0.75 to - 0.54) ^b	Responsiveness
		Men:	- 0.60 (- 0.75 to - 0.38) ^b	Responsiveness
		Women:	- 0.70 (- 0.80 to - 0.56) ^b	Responsiveness
PROMIS PI	Baseline	Total:	- 0.78 (- 0.84 to - 0.70)	Construct validity
		Men:	- 0.78 (- 0.87 to - 0.65) ^b	Construct validity
		Women:	- 0.78 (- 0.85 to - 0.68) ^b	Construct validity
	12 months	Total:	- 0.72 (- 0.82 to - 0.60) ^a	Construct validity
		Men:	– 0.71 (– 0.83 to – 0.53) ^a	Construct validity
		Women:	- 0.72 (- 0.86 to - 0.55) ^a	Construct validity
	Change	Total:	- 0.60 (- 0.70 to - 0.47) ^b	Responsiveness
		Men:	- 0.62 (- 0.77 to - 0.41) ^b	Responsiveness
		Women:	- 0.59 (- 0.72 to - 0.41) ^b	Responsiveness
PROMIS PF	Baseline	Total:	0.82 (0.74 to 0.88) ^a	Construct validity
		Men:	0.83 (0.69 to 0.91) ^a	Construct validity
		Women:	0.83 (0.75 to 0.89) ^b	Construct validity
	12 months	Total:	0.81 (0.70 to 0.88) ^a	Construct validity
		Men:	0.84 (0.74 to 0.91) ^a	Construct validity
		Women:	0.79 (0.64 to 0.89) ^a	Construct validity
	Change	Total:	0.49 (0.33 to 0.63) ^a	Responsiveness
		Men:	0.57 (0.35 to 0.74) ^b	Responsiveness
		Women:	0.44 (0.20 to 0.64) ^a	Responsiveness

Table 2 (continued)

PROMIS Patient Reported Outcomes Measurement Information System; OKS Oxford Knee Score; PAIN pain intensity; PI pain interference; PF physical function; black versus grey font color: the confidence interval of the correlation does not overlap/overlaps with the preset correlation threshold

Published online: 26 May 2023

Reference

 Stephan A, Stadelmann VA, Preiss S et al (2023) Measurement properties of PROMIS short forms for pain and function in patients receiving knee arthroplasty. J Patient Rep Outcomes 7:18. https://doi.org/10.1186/ s41687-023-00559-x

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

^a Spearman's rank correlation coefficient, r_s

^b Pearson's correlation coefficient, r